

Pediatrics QuickREF for 2025 Learning Objectives

Neck Mass—Learning Objective 29 > Quick Points

Page 242

Text currently reads:

- Infectious/Inflammatory masses
 - Reactive lymphadenopathy (LA): soft, discrete, mobile, minimally tender; may or may not be inflamed; often referred to as “shotty”
 - Infectious lymphadenitis: isolated, may be asymmetric, tender, warm, erythematous, sometimes fluctuant; less mobile and discrete than reactive lymphadenopathy
 - Viral (eg, EBV, CMV, **toxoplasmosis**, adenovirus, enterovirus, HSV, measles, HIV)
 - Typically bilateral, acute LA
 - Bacterial (eg, *Streptococcus*, *Staphylococcus*, tuberculosis)
 - Unilateral acute LA: most commonly due to group A *Streptococcus* or *Staphylococcus aureus*, generally respond to oral antibiotics (eg, amoxicillin/clavulanate, clindamycin)
 - Unilateral subacute/chronic LA: nontuberculous mycobacteria, *Bartonella henselae* (cat-scratch disease), tuberculosis
 - Mycobacterial (tuberculosis, atypical mycobacteria)

Text should read:

- Infectious/Inflammatory masses
 - Reactive lymphadenopathy (LA): soft, discrete, mobile, minimally tender; may or may not be inflamed; often referred to as “shotty”
 - Infectious lymphadenitis: isolated, may be asymmetric, tender, warm, erythematous, sometimes fluctuant; less mobile and discrete than reactive lymphadenopathy
 - Viral (eg, EBV, CMV, ~~toxoplasmosis~~, adenovirus, enterovirus, HSV, measles, HIV)
 - Typically bilateral, acute LA
 - Bacterial (eg, *Streptococcus*, *Staphylococcus*, tuberculosis)
 - Unilateral acute LA: most commonly due to group A *Streptococcus* or *Staphylococcus aureus*, generally respond to oral antibiotics (eg, amoxicillin/clavulanate, clindamycin)
 - Unilateral subacute/chronic LA: nontuberculous mycobacteria, *Bartonella henselae* (cat-scratch disease), tuberculosis
 - Mycobacterial (tuberculosis, atypical mycobacteria)