

20th Edition Internal Medicine Core

Dermatology:

Page 3-1, Common Skin Problems > Atopic Dermatitis (AD)

<i>Text currently reads:</i>	<i>Text should read:</i>
<i>Staphylococcus aureus</i> infection, appearing as honeycombing and weeping, is often present in severe cases.	<i>Staphylococcus aureus</i> infection, appearing as honey-colored crust and weeping, is often present in severe cases.

Page 3-15, Inflammatory Skin Disorders > Vasculitis

<i>Text currently reads:</i>	<i>Text should read:</i>
Palpable purpura is the extravasation of red blood cells into the skin and is commonly caused by a small vessel vasculitis . Skin biopsy typically displays leukocytoclastic vasculitis.	Palpable purpura is the extravasation of red blood cells into the skin. It is palpable because of the inflammatory milieu surrounding small vessels . Skin biopsy typically displays leukocytoclastic vasculitis, which consists of both neutrophils and fibrin deposition .

Page 3-37, HIV- and AIDS-Related Skin Lesions

<i>Text currently reads:</i>	<i>Text should read:</i>
– Kaposi sarcoma in HIV is frequently associated with HHV-8.	– Kaposi sarcoma in HIV is frequently caused by HHV-8.

Infectious Disease:

Page 4-52, Organism-Based Review > Bacteria > Gram-Negative Bacteria (GNB) > *Brucella*

<i>Text currently reads:</i>	<i>Text should read:</i>
Resistance to a single agent is common, so treatment requires 1 of the following regimens: 1) Doxycycline for 6 weeks + streptomycin for 14–21 days or gentamicin for 7–10 days 2) Doxycycline + rifampin × 6–8 weeks 3) Avoid doxycycline in pregnant persons and give rifampin + doxycycline × 6 weeks.	Resistance to a single agent is common, so treatment requires 1 of the following regimens: 1) Doxycycline for 6 weeks + streptomycin for 14–21 days or gentamicin for 7–10 days 2) Doxycycline + rifampin × 6–8 weeks 3) Avoid doxycycline in pregnant persons. Before 36 weeks gestation give rifampin and trimethoprim-sulfamethoxazole (TMP-SMX) for 6 weeks (supplement with folic acid when using TMP-SMX). Give rifampin alone at ≥ 36 weeks gestation and continue until delivery.

Nephrology:

Page 7-4, Renal Tests > Estimating Kidney Function > Glomerular Filtration Rate (GFR)

<i>Text currently reads:</i>	<i>Text should read:</i>
<p>The Cockcroft-Gault formula is another acceptable way to estimate GFR. Although it is less accurate than the MDRD and CKD-EPI equations, it can be calculated from basic variables with just the calculator on your smartphone:</p> $\text{CrCl} = \frac{(140 - \text{age}) \times (\text{weight}) \times (0.85 \text{ if female})}{72 \times \text{S}_{\text{Cr}}}$ <p style="text-align: right;">(Eq. 1)</p>	<p>The Cockcroft-Gault formula is another acceptable way to estimate GFR. Although it is less accurate than the MDRD and CKD-EPI equations, it can be calculated from basic variables with just the calculator on your smartphone:</p> $\text{CrCl} = \frac{(140 - \text{age}) \times (\text{weight}) \times (0.85 \text{ if female})}{72 \times \text{S}_{\text{Cr}}}$ <p style="text-align: right;">(Eq. 1)</p>

Page 7-37, Acid-Base Disorders > Acid-Base Balance

<i>Text currently reads:</i>	<i>Text should read:</i>
$\text{pH} = 6.10 + \log \left(\frac{\text{HCO}_3^-}{0.03 \times \text{pCO}_2} \right)$	$\text{pH} = 6.10 + \log \left[\frac{\text{HCO}_3^-}{0.03 \times \text{pCO}_2} \right]$

Page 7-48, Disorders of Water Balance > Hyponatremia > Euvolemic Hyponatremia > Primary Polydipsia

<i>Text currently reads:</i>	<i>Text should read:</i>
<p>Primary polydipsia is a psychiatric disorder in which patients consume large amounts of water (as much as v).</p>	<p>Primary polydipsia is a psychiatric disorder in which patients consume large amounts of water (as much as 1 L/hour).</p>

Oncology:

Page 9-36, Hypercalcemia of Malignancy

<i>Text currently reads:</i>	<i>Text should read:</i>
$\text{Ca}^{2+}_{\text{corrected}} = \text{Ca}^{2+}_{\text{measured}} + 0.8 \times (4.0 - \text{alb}_{\text{measured}})$	$\text{Ca}^{2+}_{\text{corrected}} = \text{Ca}^{2+}_{\text{measured}} + [0.8 \times (4.0 - \text{alb}_{\text{measured}})]$